

# SupaWetter

Reg.No. 16-ADJV-2-3


**A 100% non-ionic organosilicone-based wetter / spreader / penetrant for use with agricultural remedies to reduce the surface tension of the spray solution**

## ACTIVE INGREDIENT

Polyether –polymethylsiloxane 1000 g/l  
Inert Ingredient 0 g/l

## CHEMICAL GROUP

Organo Silicone



**HARMFUL IF SWALLOWED**



# Intralab

Manufactured by:

**INTROLAB (PTY) LTD** Reg No.: 2005/039779/07  
12 Oudepont street, Wellington Industrial,  
Wellington, South Africa, 7655  
Tel: +27 (021) 864 2145/6 • Email: info@intralab.co.za

[www.intralab.co.za](http://www.intralab.co.za)

Distributed by:

**INTROLAB ZIMBABWE**  
19 Greendale Ave, Greendale,  
Harare, Zimbabwe  
Tel: +263 71 263 5535

17017 01/2017 SupaWetter

See container side for batch number, manufacture date, expiry date and content.



## Adjuvant Range

SEE ENCLOSED  
FOR FULL DETAILS

## TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

### PRECAUTIONS:

1. Handle with care. Avoid any contact with the wetter.
2. WEAR PROTECTIVE CLOTHING i.e. gloves and overalls, rubber boots when handling this wetter.
3. **DO NOT** eat, drink or smoke while applying this pesticide.
4. Remove clothing on completion of spraying and wash hands and face thoroughly with soap and water.
5. DANGEROUS TO ANIMALS, BIRDS AND BEES.
6. **DO NOT** CONTAMINATE DRINKING POOLS, DAMS, RIVERS AND WATERWAYS.
7. KEEP APART FROM FOOD AND FOODSTUFFS.
8. Dispose empty containers by triple rinsing and immediately emptying washings into the spray tank for spraying onto the crop. Perforate and flatten rinsed container to make it unusable for further use. Bury the crushed container in a disposal pit in a secure location away from water sources or arable land. **DO NOT** use container for any other purpose.
9. Store in original containers and **KEEP UNDER LOCK AND KEY.**

**SYMPTOMS OF POISONING:** Causes moderate irritation to the eyes and skin. Inhalation may result in irritation of respiratory and tract. Ingestion may result in gastric distress and vomiting.

### First aid treatment:

**Ingestion:** Have the victim rinse mouth thoroughly with water. Administer water to dilute the material and if vomiting occurs have the victim lean forward with head down to avoid breathing in of vomits. Rinse the mouth and continue to administer water. **DO NOT** induce vomiting.

**Skin contact:** Remove any contaminated clothing. Wash the skin thoroughly with soap and water. **DO NOT** rub skin hard and wash all contaminated clothing before re-use.

**Eye contact:** Immediately flush eyes with plenty of clean water for at least 10 - 15 minutes.

**Inhalation:** Remove person to fresh air. Give oxygen if breathing is difficult.

**Notes to physician:**  
Treat symptomatically

**WARRANTY:** The excellent results which this product normally yields are not however warranted or guaranteed as the effectiveness depends on weather and soil conditions, application techniques and other factors outside the control of the manufacturers and the suppliers. No warranty whatsoever express or implied, is therefore given concerning the effectiveness of this product and responsibility is specifically excluded for any damage injury or loss of any kind during or resulting from its handling, use or storage whether or not such has been in accordance with directions. This product is warranted to consist of the ingredients shown on the label.

**Compatibility:** Supawetter is a new generation surfactant based on organo-modified siloxane technology. Supawetter is a highly effective non-ionic surfactant / wetter / spreader / penetrant. The Supawetter effectiveness may allow the use of the lowest labeled pesticide rate.

**SHELF LIFE:** A minimum of 2 years from date of manufacture if kept under constant cool and dry conditions.

DIRECTIONS FOR USE: (Use only as indicated)

Use with	Dosage rate	Remarks
<b>HERBICIDES</b>	150 – 250 ml/ha	Use the higher rate for difficult- to- wet weeds
<b>INSECTICIDES &amp;</b>	100 – 250 ml/ha	Row crops (Cereals)
<b>FUNGICIDES</b>	250 ml/ha	Pome fruit, stone fruit, vines
	250 ml/ha	Sub-tropical fruit
	3 -5 ml/100 ℓ water	Citrus
	250 ml/ha	Horticultural crops and ornamentals
<b>Plant growth regulators</b>	25 – 50 ml/100 ℓ water	Avoid run-off. Do not exceed 500 ml Supawetter per hectare.
<b>Defoliant</b>	100 – 200ml/ha	
<b>Foliar applied micro nutrients</b>	100 – 200ml/ha	Row crops (Cereals)
	250 ml/ha	Pome fruit, stone fruit, vines
	250 ml/ha	Sub-tropical fruit
	3 -5 ml/100 ℓ water	Citrus
	250 ml/ha	Horticultural crops and ornamentals
<b>Apple Calcium</b>	100 ml/ha	Apples
<b>Nitrate applications</b>		
	1 – 1,5 ℓ/ha	As a soil-wetting agent. Apply immediately prior to irrigation either with a boom sprayer or via the irrigation system. Repeat
<b>Golf &amp; Turf</b>		at 14-day intervals
	30 ml/100m <sup>2</sup>	Spot treatment of hydrophobic areas. Irrigate immediately after application

MIXING INSTRUCTIONS:

- Fill the spray tank 2/3 full of water. Commence agitation. If foaming is anticipated, add **Anti-Foam** before pesticide and **SupaWetter**.
- Add pesticide or Foliar fertilizers as directed by their specific label.
- Continue agitation.
- Add **SupaWetter** last to the spray tank and continue filling while maintaining mild agitation.