

### Product Description:

**IntroSpoor** is a water soluble powder containing micro-elements for use as soil or foliar application to prevent or correct deficiencies in plants.

### Compatibility:

- **IntroSpoor** is compatible with most agrochemicals.
- It is however advisable to do a miscibility test prior to mixing this product with other agrochemicals.
- Always follow the standard tank mixing procedure.

### Storage:

- Store in the original, well-sealed packaging away from direct sunlight in a cool dry environment.
- Keep away from children, uninformed people, animals and foodstuffs.

### Warning:

- Handle with care.
- Do not re-use the packaging for any other purpose.
- Only use where there is a recognized need.

### Application instructions:

- **IntroSpoor** should preferably be applied in the early morning or late afternoon.
- Do not apply to plants that are undergoing moisture or heat stress.
- **Foliar application:** Apply in sufficient water to ensure good even coverage of the foliage.
- **Soil application:** Apply in sufficient water to ensure an even distribution. Incorporate or wash into the root area immediately after application.

**Manufactured & Registered by:**  
**INTROLAB CHEMICALS (PTY) LTD**  
(Reg. No. 2005/039779/07)  
12 Oudepont Street, Wellington  
Industrial, Wellington,  
South Africa, 7655  
Tel: +27 21 864 2145/6  
Email: [info@introlab.co.za](mailto:info@introlab.co.za)



# INTROSPOR

**IntroSpoor** is a water soluble powder containing micro-elements for use as soil or foliar application to prevent or correct deficiencies in plants.

Reg. No. B5809  
Act 36 of 1947, Fertilizer Group 2

### ACTIVE INGREDIENTS:

Zinc (Zn)*	37.5 g/kg
Iron (Fe)*	36 g/kg
Boron (B)	35 g/kg
Manganese (Mn)*	32.5 g/kg
Copper (Cu)*	2.25 g/kg
Molybdenum (Mo)	3 g/kg
*EDTA Chelated	

**WARRANTY:** Although this product has been extensively tested under large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application.

**Batch No.:**

PACKING  
SIZE **5kg**

### DIRECTIONS FOR USE (USE ONLY AS DIRECTED)

#### FOLIAR APPLICATION

CROP	Grain crops	Vegetables
RATE/HA	1 – 3 kg	0.5 – 3 kg
<b>Remarks:</b> Apply as soon as there is enough leaf material intercept spray. Repeat at 10-14 day intervals if necessary. Do not exceed 6kg/ha per season.		

CROP	Subtropical fruits
RATE/HA	2 – 3 kg
<b>Remarks:</b> Apply post-harvest, before flowering and repeat at 80% petal fall if necessary. Do not exceed 6kg/ha per season.	

CROP	Nut trees
RATE/HA	2 – 3 kg
<b>Remarks:</b> Apply post-harvest, before flowering. Repeat at 80% petal fall if necessary. Apply in 2000-3000 L water per hectare. Do not exceed 6kg/ha per season.	

CROP	Stone & pome fruit
RATE/HA	2 – 3 kg
<b>Remarks:</b> Apply post-harvest (6-8 weeks before leaf drop). Repeat at bloom to 80% petal drop. Apply a third application during pit hardening if necessary. Do not exceed 6kg/ha per season. Apply in 500-1500 L water per hectare.	

CROP	Pastures
RATE/HA	1 – 3 kg
<b>Remarks:</b> Apply after each cut or grazing when there is enough leaf material to absorb spray (typically a plant height of 10-15cm). Ensure 14 days between application and grazing. Do not exceed 6kg/ha per season.	

#### FERTIGATION

CROP	Pot plants
DOSAGE RATE/1000L WATER	10 – 20g
<b>Remarks:</b> Apply by means of irrigation water or apply to the drip zone by hand. Use at least 5 – 10 mm water to wash the product into root zone.	

CROP	Seedlings in trays
DOSAGE RATE/1000L WATER	10 – 20g
<b>Remarks:</b> Apply 2-3 times from emergence to transplant in 1000L water.	

CROP	Fruit crops
DOSAGE RATE/1000L WATER	2 kg/ha
<b>Remarks:</b> Apply by means of irrigation water or apply to the drip zone by hand. Use at least 5 – 10 mm water to wash the product into root zone.	

Consult with a qualified Agronomist or Agricultural Advisor for the correct dosage and application method of the product.