

INTRO CAL



Reg. No.: B5715
(Act 36 of 1947)
Fertilizer Group 2

Intro Cal is an Amino Acid chelated Calcium containing liquid fertilizer, specifically designed for use as a foliar feed to prevent and correct Calcium deficiencies.

Active Ingredient:

- | | | | |
|-------------|------|---------|----------|
| * Calcium | (Ca) | 80 g/l | 64 g/kg |
| Nitrogen | (N) | 48 g/l | 38 g/kg |
| Amino Acids | | 170 g/l | 136 g/kg |
- * Amino Acid and Carbohydrate Chelated
SG (@ 20 °C) = 1.25 ± 0.02



Registered, Manufactured & Distributed by:

INTROLAB (PTY) LTD
(Reg. No. 2005/039779/07)

12 Oudepont Street
Wellington Industrial
Wellington
South Africa
7655
Tel: +27 21 864 2145/6
Email: info@introlab.co.za



Introlab
outperform the norm

Version: April 2021

www.introlab.co.za

See enclosed for full details. ➡

See container side for both Batch No. and Manufacture Date. Product expires 2 years after date of manufacture.

Product Description:

- **Intro Cal** is an amino acid complexed Calcium containing liquid fertilizer.
- The amino acid complexed nutrient will enhance absorption by plants.
- **Intro Cal** is suitable for foliar application.
- **Intro Cal** is readily taken up by plants.

Compatibility:

- **Intro Cal** is compatible with most agricultural remedies.
- It is advisable to do a miscibility test prior to mixing **Intro Cal** with other agricultural products

Intro Cal could lower the spray water pH.

The use of a pH buffer such as, AquaTech is recommended when tank mixing **Intro Cal** with other Agrochemicals that are susceptible to low pH conditions.

Do not apply **Intro Cal**:

- In a tank mix with Copper containing fungicides.
- Onto plants with Copper fungicide containing residues.
- Within 30 days after Copper containing fungicide applications.
- In a tank mix with products that contain high levels of phosphate or phosphonate.

Contact the Manufacturer or Crop advisor before tank mixing **Intro Cal** with other Agrochemicals.

Mixing Instructions:

- Shake the container thoroughly before use.
- Follow the standard tank mixing procedure if this product is applied in combination with other agricultural products.
- Ensure good agitation throughout application.

Storage:

- Store in the original, well-sealed container away from direct sunlight in a cool dry environment.
- Keep away from children, uninformed people, animals and foodstuffs.

Application Instructions:

- **Intro Cal** should preferably be applied in the early morning or late afternoon.
- Do not apply to plants that are undergoing moisture or heat stress.
- Ensure good even coverage of the plants.
- For optimum results **Intro Cal** should be used in a balanced plant nutrition programme.
- Consult your chemical distributor, consultant or the registration holder should crop specific programmes be required, or if any additional information and guidance is required concerning the use of this product.

Warning:

- Handle with care.
- Do not re-use the container for any other purpose.

Intro Cal could lower the spray water pH.

The use of a pH buffer such as, AquaTech is recommended when tank mixing **Intro Cal** with other Agrochemicals that are susceptible to low pH conditions.

Do not apply Intro Cal:

- In a tank mix with Copper containing fungicides.
- Onto plants with Copper fungicide containing residues.
- Within 30 days after Copper containing fungicide applications.
- In a tank mix with products that contain high levels of phosphate or phosphonate.

Contact the Manufacturer or Crop advisor before tank mixing **Intro Cal** with other Agrochemicals.

Directions for use: (Use only as directed)

Use the lower rates as maintenance and the higher rate to correct deficiencies:

Crop	Rate/ha	Remarks
Berries: i. Blueberries, Blackberries, Raspberries	2 - 7 ℓ 2 - 7 ℓ	i. Commence at flowering ii. Every second week until the end of harvest. Apply in 300 ℓ water/ha
ii. Strawberries	2 - 7 ℓ 2 - 7 ℓ	i. Two weeks after transplant ii. Every second week until the end of harvest. Apply in 200 ℓ water/ha
Bulb, Tuber & Root Crops: i. Onions, Garlic	2 - 3 ℓ 2 - 3 ℓ 2 - 3 ℓ	ii. 4 Weeks post emergence. ii. 6 Weeks post emergence iii. 8 Weeks post emergence Apply in 400 liter water/ha
ii. Potatoes, Sweet Potatoes	4 - 5 ℓ 4 - 5 ℓ 4 - 5 ℓ 4 - 5 ℓ	i. Apply at first emergence ii. Apply 4 Weeks post emergence iii. Weeks post emergence iv. 12 Weeks post emergence Apply in 200 - 400 ℓ water/ha
iii. Beetroot, Carrots	3 - 4 ℓ 3 - 4 ℓ 3 - 4 ℓ	i. 2-3 Weeks post emergence ii. 14 days later iii. 14 days later Apply in 200-300 ℓ water/ha
Cruciferae: Broccoli, Cabbage, Brussel Sprouts, Cauliflower	3 - 4 ℓ 3 - 4 ℓ 3 - 4 ℓ	i. 2-3 Weeks post-transplant ii. 14 days later iii. 14 days later Apply in 200-300 ℓ water/ha
Cucurbits: Cucumbers, Marrows, Melons, Pumpkins, Baby marrows, Watermelons	2 - 4 ℓ 2 - 4 ℓ 2 - 4 ℓ	i. Directly after fruit set ii. 10 days later iii. 10 days later Apply in 200 ℓ water / ha

Directions for use: (Use only as directed)

Crop	Rate/ha	Remarks
Grain Crops: i. Maize, Sorghum	3 – 4 ℓ	i. Apply at tasseling stage via overhead irrigation or tractor application in 200 ℓ water/ha.
ii. Barley, Oats, Wheat	2 – 3 ℓ	i. Apply at tillering ii. 14 days later Apply in 200 ℓ of water/ha.
Grapes: Table & Wine grapes	5 - 10 ℓ 5 - 10 ℓ 5 - 10 ℓ	i. Apply directly after fruit set ii. Apply 14 days later iii. Apply 14 days later Apply in 500 - 1000 ℓ of water/ha.
Leafy vegetables: Lettuce, Spinach	2 – 3 ℓ	i. Apply 3 weeks post emergence, repeat application every 14 days until one month before harvest. Apply in 200 ℓ of water/ha.
Legumes: Dry Beans, Peas, Other Beans	3 – 5 ℓ 3 – 5 ℓ	i. Start application on onset of flowering ii. 14 days later Apply in 200 ℓ of water/ha.
Mediterranean Fruit: Figs, Olives, Pomegranates	5 – 10 ℓ 5 – 10 ℓ	i. Apply at petal fall ii. 14 days later Apply in 750 – 1500 ℓ water/ha.
Nuts: i. Almonds	5 – 10 ℓ 5 – 10 ℓ 5 – 10 ℓ	i. 75% Petal fall ii. Two weeks later iii. Two weeks later Apply in 1000 – 2000 ℓ water/ha.

Directions for use: (Use only as directed)

Crop	Rate/ha	Remarks
ii. Macadamia	5 – 10 ℓ 5 – 10 ℓ	i. August at full bloom ii. 30 Days later Apply in 1500 -3000 ℓ water/ha.
iii. Pecan, Wall Nuts	5 – 10 ℓ 5 – 10 ℓ	i. 2 Weeks post parachute stage (Middle October) ii. Middle December Apply in 1000 – 2000 ℓ water/ha.
Oil Seed Crops:		
i. Groundnuts	3 – 5 ℓ 3 – 5 ℓ	i. Start application at flowering ii. Repeat 14 days later Apply in 200 ℓ of water/ha.
ii. Soy Beans	3 – 5 ℓ 3 – 5 ℓ	i. Start application on onset of flowering ii. 14 days later Apply in 200 ℓ of water/ha.
iii. Sunflower	3 – 5 ℓ	i. Apply just before flowering in 200 ℓ water
Pineapples	3 – 5 ℓ 3 – 5 ℓ 3 – 5 ℓ	i. Apply at blue flower stage (Middle February) ii. Dead Flower Stage (Early March) iii. 2 weeks later Apply in 500 ℓ water/ha.
Pome fruit:		
Apples, Pears, Quinces	5 – 10 ℓ 5 – 10 ℓ	i. Apply from petal fall ii. Repeat application every 10-14 days until one month before harvest Apply in 1000 – 3000 ℓ water /ha according to Tree Row Volume.
Solanaceous Crops:		
i. Chillies, Paprika, Peppers, Tomatoes :	3 – 5 ℓ 3 – 5 ℓ	i. Apply at the end of flowering on first truss ii. Repeat application every 7-10 days until one month before harvest Apply in 400 ℓ water/ha.
ii. Tobacco	3 – 5 ℓ	i. Apply as a full cover spray 3 weeks after transplant in 500 ℓ water/ha.

Directions for use: (Use only as directed)

Crop	Rate/ha	Remarks
Stone Fruit: Apricots, Nectarines, Peaches, Plums	5 – 10 ℓ 5 – 10 ℓ	i. Apply from petal fall ii. Repeat application every 10-14 days until one month before harvest Apply in 1000 – 3000 ℓ water /ha according to Tree Row Volume.
Subtropical Fruit: i. Avocados	5 – 10 ℓ 5 – 10 ℓ 5 – 10 ℓ	i. Cauliflower stage ii. Full bloom iii. Fruit set Direct sprays at the flower trusses. Apply in 3000 – 5000 ℓ water/ha.
ii. Guavas	3 – 6 ℓ 3 – 6 ℓ	i. Apply at fruit set. ii. Repeat application every 14 days until one month before harvest. Apply in 1500 – 3000 ℓ water/ha.
iii. Lychees, Mangoes	10 – 15 ℓ 10 – 15 ℓ	i. Apply at fruit set ii. Repeat application 14 days later Direct sprays at the flower trusses. Apply in 2000 liter water/ha.
iv. Citrus	10 -15 ℓ 10 -15 ℓ	i. Apply at 50-100% petal fall (September) ii. 2-4 weeks post 100% petal fall Apply according to Tree Row Volume or 1000 – 4000 ℓ water/ha.
Other Crops	3 – 5 ℓ	Start application from early fruit development stage and repeat applications if necessary at 10-14 day intervals. Apply in a minimum of 500 ℓ water/ha.

Warranty:

Although this product has been extensively tested under large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application